

PRODUCE:

ΡΟΤΑΤΟ

ТҮРЕ	Washed	VARIETY	Gold Star
CLASS	One	NOTES	
GENERAL APPEARANCE CRITERIA			
Colour	Light golden skin which may be mottled to lightly russetted; yellowish to creamy flesh.		
Visual Appearance	Shallow eyes, moderately pronounced lenticels; free from foreign matter; skinning is allowable for new seasons tubers		
Sensory	Firm when tested with finger pressure, not soft or spongy; free from foreign tastes or odours; no 'off' or foreign odours or tastes.		
Shape	Uniform round to oval; nil with deformed shape.		
Size	Weight range 60 - 110 g packs < 1 kg; 110 - 350 g > 1kg packs.		
MAJOR DEFECTS			
Insects	With obvious live insects (eg potato moth, potato wireworm)		
Diseases	With fungal or bacterial rots (eg pink rot, dry rot, gangrene, late blight or bacterial wilt).		
Physical/Pest Damage	With numerous raised or sunken corky growths on the potato skin (powdery scab, common scab). With evidence of insect, rodent, or nematode damage. With cuts or splits >10mm long, and/or >1mm wide and/or >3 mm deep. With severely malformed or root constricted tubers, or tubers with advanced secondary growth. With green areas on the skin or an overall green tinge (exposed to light, toxic compounds may form) With deep, dark bruising and softened flesh.		
Physiological Disorders	With sprouting from the eyelets (overstored tubers). With dark pink or black areas at the centre of the tuber (black heart, due to low oxygen). With severe brown spotting of the flesh (brown fleck) or hollow cavities inside the tubers (hollow heart).		
Temperature Injury	With obvious shrivelling and softened, spongy flesh (dehydration). With reddish brown blotches or soft, water-soaked areas in the flesh (freezing injury).		
MINOR DEFECTS			
Physical/Pest Damage	With cuts or splits <10mm long, <1mm wide and <3 mm deep (not removable by one pass peeling). With superficial healed bruises, scuffs, pressure or rub marks affecting in total >1 sq cm and >3 mm deep.		
Skin Marks/Blemishes	With moderate development of a silvery sheen, brown markings or skin cracking affecting >2 sq cm. With moderate black speckling affecting in aggregate >1 sq cm of surface area.		
Physiological Disorder	With enlarged, swollen lenticels (>1mm diameter) affecting >3 sq cm. With healed growth cracks or splits >2mm wide or >2mm deep (not removable by one pass peeling).		
CONSIGNMENT CRITERIA			
Tolerance Per Consignment		limit) to be < 2 defects per item Total m . Total major defects must not exceed 2	
Packaging & Labelling	Packaging manufactured from new food grade materials or sanitised returnable crates. All labelling must meet the current legislative requirements. Labelling to identify grower's name/brand (plus growers name/code if via a packhouse), address, contents, class, size and/or minimum net weight. Produce to identify Country of Origin (eg. Produce of Australia) on outer container.		
Shelf Life	Produce must provide not less than 14 days clear shelf life from date of receival.		
Receival Conditions	Compliance with Quarantine Treatments (if required) for Interstate Consignment. Stacked onto a stabilised pallet as pre-ordered. Refrigerated van with air bag suspension, unless otherwise approved. Pulp Temperature 12 - 20°C		
Chemical & Containment Residues	All chemicals used pre/postharvest must be registered and approved for use in accordance with the requirements of the APVMA regulatory system. Residues, Contaminants and Heavy Metals to comply with the FSANZ Food Standards Code MRL's and ML's.		
Food Safety Requirements	Produce is to be grown and packed under a HACCP based food safety program that is subject to an annual third- party audit. A copy of current certification to be forwarded to receiver.		

*Specifications reviewable: e.g. to account for specific regional effects or adverse seasonal impacts on quality or early or late seasonal variances as agreed and communicated formally in writing.