Effective: 1 November 2006



PRODUCE: POTATO

ТҮРЕ	Washed	VARIETY	Roseval
CLASS	One	NOTES	

Colour	Dark red skin with uniform clean, bright bloom; yellow flesh.		
/isual Appearance	Shallow to medium depth eyes; free from foreign matter; skinning is allowable for new seasons tubers.		
Sensory	Firm when tested with finger pressure, not soft or spongy; free from foreign tastes or odours; no 'off' or foreign odours or tastes.		
Shape	Approximately long to oblong.		
Size	Weight range 60 - 110 g packs < 1 kg; 110 - 350 g > 1kg packs		
MAJOR DEFECTS			
nsects	With obvious live insects (eg potato moth, potato wireworm)		
Diseases	With fungal or bacterial rots (eg pink rot, dry rot, gangrene, late blight or bacterial wilt).		
Physical/Pest Damage	With numerous raised or sunken corky growths on the potato skin (powdery scab, common scab). With evidence of insect, rodent, or nematode damage. With cuts or splits >20mm long, and/or >2mm wide and/or >3 mm deep. With severely malformed or root constricted tubers, or tubers with advanced secondary growth. With deep, dabruising and softened flesh. With green areas on the skin or an overall green tinge (exposed to light, toxic compounds may form)		
Physiological Disorders	With sprouting from the eyelets (overstored tubers). With severe brown spotting of the flesh (brown fleck) or hollow cavities inside the tubers (hollow heart). With dark pink or black areas at the centre of the tuber (black heart, due to low oxygen).		
Femperature Injury	With reddish brown blotches or soft, water-soaked areas in the flesh (freezing injury). With obvious shrivelling and softened, spongy flesh (dehydration).		
MINOR DEFECTS			
Physical/Pest Damage	With cuts or splits <20mm long, <2mm wide and <3 mm deep (not removable by one pass peeling). With superficial healed bruises, scuffs, pressure or rub marks affecting in total >2 sq cm and >3 mm deep.		
Skin Marks/Blemishes	With moderate development of a silvery sheen, brown markings or skin cracking affecting >5 sq cm. With moderate black speckling affecting in aggregate >2 sq cm of surface area.		
Physiological Disorder	With healed growth cracks or splits >3mm wide or >3mm deep (not removable by one pass peeling). With enlarged, swollen lenticels (>1mm diameter) affecting >5 sq cm.		
CONSIGNMENT CRITERIA			
Folerance Per Consignment	Total minor defects (within allowance limit) to be < 2 defects per item Total minor defects (outside allowance limust not exceed 10% of consignment. Total major defects must not exceed 2% of consignment. Combined Tot not to exceed 10%.		
Packaging & Labelling	Packaging manufactured from new food grade materials or sanitised returnable crates. All labelling must meet current legislative requirements. Labelling to identify grower's name/brand (plus growers name/code if via a packhouse), address, contents, class, size and/or minimum net weight. Produce to identify Country of Origin (exproduce of Australia) on outer container.		
Shelf Life	Produce must provide not less than 14 days clear shelf life from date of receival.		
Receival Conditions	Compliance with Quarantine Treatments (if required) for Interstate Consignment. Stacked onto a stabilised pallet as pre-ordered. Refrigerated van with air bag suspension, unless otherwise approved. Pulp Temperature 12 - 20°C		
	All chemicals used pre/postharvest must be registered and approved for use in accordance with the requirements of the APVMA regulatory system. Residues, Contaminants and Heavy Metals to comply with the FSANZ Food Standards Code MRL's and ML's.		
Chemical & Containment Residues			